

Portmanteau Test Statistics

Esam Mahdi

University of Western Ontario

A. Ian McLeod

University of Western Ontario

Abstract

In this vignette, we give a brief description about the portmanteau test statistics given in the **portes** package. Some applications, including two examples from [Mahdi and McLeod \(2011\)](#) are given in this vignette as well.

Keywords: ARMA models, Monte-Carlo significance test, Portmanteau test, VARMA models .

1. Box and Pierce portmanteau test

In the univariate time series, [Box and Pierce \(1970\)](#) introduced the portmanteau statistic

$$Q_m = n \sum_{\ell=1}^m \hat{r}_\ell^2 \quad (1)$$

where $\hat{r}_\ell = \sum_{t=\ell+1}^n \hat{a}_t \hat{a}_{t-\ell} / \sum_{t=1}^n \hat{a}_t^2$, and $\hat{a}_1, \dots, \hat{a}_n$ are the residuals. This test statistic is implemented in the R function `BoxPierce()` and can be used in the multivariate case as well. It has a chi-square distribution with $k^2(m - (p + q))$ degrees of freedom where k represents the dimension of the time series. The usage of this function is extremely simple:

```
BoxPierce(obj, lags = seq(5, 30, 5), order = 0, SquaredQ = FALSE),
```

where `obj` is a univariate or multivariate series with class `"numeric"`, `"matrix"`, `"ts"`, or `("mts" "ts")`. It can be also an object of fitted time-series model with class `"ar"`, `"arima0"`, `"Arima"`, `"varest"`, `"FitAR"`, or `"FitFGN"`. `lags` is a vector of numeric integers represents the lag values, m , at which we need to check the adequacy of the fitted model. The argument `order` is used for degrees of freedom of asymptotic chi-square distribution. If `obj` is a fitted time-series model with class `"ar"`, `"arima0"`, `"Arima"`, `"varest"`, `"FitAR"`, or `"FitFGN"` then no need to enter the value of `order` as it will be automatically determined. In general `order = p + q`, where `p` and `q` are the orders of the autoregressive (or vector autoregressive) and moving average (or vector moving average) models respectively. `order = 0` is used for testing random series, fractional gaussian noise, or generalized autoregressive conditional heteroscedasticity. Finally, when `SquaredQ = TRUE`, then apply the test on the squared values. This checks for Autoregressive Conditional Heteroscedastic, ARCH, effects. When `SquaredQ = FALSE`, then apply the test on the usual residuals.

Note that the function `portest()` with the arguments `test = "BoxPierce"`, `MonteCarlo = FALSE`, and `order = 0` will gives the same results of the function `BoxPierce()`. The Monte-Carlo version of this test statistic is implemented in the function `portest()` as an argument `test = "BoxPierce"` provided that `MonteCarlo = TRUE` is selected.

```
portest(obj, lags = seq(5, 30, 5), order = 0, test = "BoxPierce", MonteCarlo = TRUE,
        nslaves = 1, NREP = 1000, InfiniteVarianceQ = FALSE, SquaredQ = FALSE)
```

1.1. Example 1

First a simple univariate example is provided. We fit an AR(2) model to the logarithms of Canadian lynx trappings from 1821 to 1934. Data is available from the R package **datasets** under the name **lynx**. This model was selected using the BIC criterion. The asymptotic distribution and the Monte-Carlo version of Q_m statistic are given in the following R code for lags $m = 5, 10, 15, 20, 25, 30$ with **snow** package using PC with two CPU's.

```
R> library("portes")
R> library("snow")
R> nslaves <- 2
R> lynxData <- log(lynx)
R> p <- SelectModel(lynxData, ARModel = "AR", Criterion = "BIC",
+   Best = 1)
R> Fitlynx <- FitAR(lynxData, p, ARModel = "AR")
R> BoxPierce(Fitlynx)
```

Lags	Statistic	df	p-value
5	6.748225	3	0.08037069
10	15.856081	8	0.04448698
15	22.631444	13	0.04631764
20	30.304179	18	0.03459211
25	34.157210	23	0.06291892
30	37.963103	28	0.09909886

```
R> portest(Fitlynx, test = "BoxPierce", nslaves = nslaves)
```

2 slaves are spawned successfully. 0 failed.

Lags	Statistic	df	p-value
5	6.748225	3	0.08491508
10	15.856081	8	0.03296703
15	22.631444	13	0.02597403
20	30.304179	18	0.02197802
25	34.157210	23	0.03296703
30	37.963103	28	0.04395604

For lags $m \geq 10$, the Monte-Carlo version of Box and Pierce test is more decisively suggests model inadequacy, whereas the asymptotic chi-square suggests inadequacy at lags 10 to 20 and adequacy otherwise. Fitting a subset autoregressive using the BIC ([McLeod and Zhang 2008](#)), the portmanteau test based on both methods, Monte-Carlo and asymptotic distribution suggest model adequacy.

```
R> SelectModel(log(lynx), lag.max = 15, ARModel = "ARp", Criterion = "BIC",
+   Best = 1)
```

```
[1] 1 2 4 10 11
```

```
R> FitsubsetAR <- FitARp(log(lynx), c(1, 2, 4, 10, 11))
R> BoxPierce(FitsubsetAR)
```

Lags	Statistic	df	p-value
5	2.382300	0	NA
10	4.258836	0	NA
15	6.532786	4	0.1627363
20	9.887818	9	0.3596432
25	13.258935	14	0.5062439
30	16.172499	19	0.6457394

```
R> portest(FitsubsetAR, test = "BoxPierce", nslaves = nslaves)
```

```
2 slaves are spawned successfully. 0 failed.
```

Lags	Statistic	df	p-value
5	2.382300	0	0.5224775
10	4.258836	0	0.7742258
15	6.532786	4	0.8311688
20	9.887818	9	0.7992008
25	13.258935	14	0.7852148
30	16.172499	19	0.7752248

1.2. Example 2

In this example we consider the monthly log stock returns of Intel corporation data from January 1973 to December 2003. First we apply the Q_m statistic directly on the returns using the asymptotic distribution and the Monte-Carlo significance test. The results suggest that returns data behaves like white noise series as no significant serial correlations found.

```
R> monthintel <- as.ts(monthintel)
R> BoxPierce(monthintel)
```

Lags	Statistic	df	p-value
5	4.666889	5	0.45786938
10	14.364748	10	0.15699489
15	23.120348	15	0.08161787
20	24.000123	20	0.24238680
25	29.617977	25	0.23891229
30	31.943703	30	0.37015020

```
R> portest(monthintel, test = "BoxPierce", nslaves = nslaves)
```

```
2 slaves are spawned successfully. 0 failed.
```

Lags	Statistic	df	p-value
------	-----------	----	---------

```

5  4.666889  5  0.4385614
10 14.364748 10  0.1518482
15 23.120348 15  0.0979021
20 24.000123 20  0.2157842
25 29.617977 25  0.2307692
30 31.943703 30  0.3166833

```

After that we apply the Q_m statistic on the squared returns. The results suggest that the monthly returns are not serially independent and the return series may suffers of ARCH effects.

```
R> BoxPierce(monthintel, SquaredQ = TRUE)
```

```

Lags Statistic df      p-value
 5  40.78073  5  1.039009e-07
10  49.57872 10  3.189915e-07
15  81.90133 15  3.131517e-11
20  86.50575 20  3.006796e-10
25  87.54737 25  7.161478e-09
30  88.55017 30  1.087505e-07

```

```
R> portest(monthintel, test = "BoxPierce", nslaves = nslaves, SquaredQ = TRUE)
```

```
2 slaves are spawned successfully. 0 failed.
```

```

Lags Statistic df      p-value
 5  40.78073  5  0.000999001
10  49.57872 10  0.000999001
15  81.90133 15  0.000999001
20  86.50575 20  0.000999001
25  87.54737 25  0.000999001
30  88.55017 30  0.000999001

```

2. Ljung and Box portmanteau test

Ljung and Box (1978) modified Box and Pierce (1970) test statistic by

$$\hat{Q}_m = n(n+2) \sum_{\ell=1}^m (n-\ell)^{-1} \hat{r}_\ell^2. \quad (2)$$

This test statistic is also asymptotically chi-square with degrees of freedom $k^2(m-p-q)$ and implemented in the R function `LjungBox()`,

```
LjungBox(obj, lags = seq(5, 30, 5), order = 0, SquaredQ = FALSE),
```

where the arguments of this function are described as before.

In R, the function `Box.test()` was built to compute the [Box and Pierce \(1970\)](#) and [Ljung and Box \(1978\)](#) test statistics only in the univariate case where we can not use more than one single lag value at a time. The functions `BoxPierce()` and `LjungBox()` are more general than `Box.test()` and can be used in the univariate or multivariate time series at vector of different lag values as well as they can be applied on an output object from a fitted model.

Note that the function `portest()` with the arguments `test = "LjungBox"`, `MonteCarlo = FALSE`, and `order = 0` will gives the same results of the function `LjungBox()`. The Monte-Carlo version of this test statistic is implemented in the function `portest()` as an argument `test = "LjungBox"` provided that `MonteCarlo = TRUE` is selected.

```
portest(obj, lags = seq(5, 30, 5), order = 0, test = "LjungBox", MonteCarlo = TRUE,
        nslaves = 1, NREP = 1000, InfiniteVarianceQ = FALSE, SquaredQ = FALSE)
```

2.1. Example 3

The built in R function `auto.arima()` in the package **forecast** is used to fit the best ARIMA model based on the AIC criterion to the measurements of the annual flow of the river Nile at Aswan from the years 1871 to 1970,

```
R> library("forecast")
```

```
R> FitNile <- auto.arima(Nile)
```

Then the `LjungBox` portmanteau test is applied on the residuals of the fitted model at lag values 5, 10, 15, 20, 25, and 30 which yields that the assumption of the adequacy in the fitted model is fail to reject.

```
R> LjungBox(FitNile)
```

Lags	Statistic	df	p-value
5	1.257698	3	0.7392018
10	9.705584	8	0.2863011
15	11.415751	13	0.5760319
20	12.861450	18	0.7997373
25	14.437766	23	0.9136466
30	17.395015	28	0.9403734

```
R> portest(FitNile, test = "LjungBox", nslaves = nslaves)
```

2 slaves are spawned successfully. 0 failed.

Lags	Statistic	df	p-value
5	1.257698	3	0.8521479
10	9.705584	8	0.3256743
15	11.415751	13	0.6023976
20	12.861450	18	0.8211788
25	14.437766	23	0.9200799
30	17.395015	28	0.9370629

3. Hosking portmanteau test

Hosking (1980) generalized the univariate portmanteau test statistics given in eqns. (1, 2) to the multivariate case. He suggested the modified multivariate portmanteau test statistic

$$\tilde{Q}_m = n^2 \sum_{\ell=1}^m (n - \ell)^{-1} \hat{\mathbf{r}}_\ell' (\hat{\mathbf{R}}_0^{-1} \otimes \hat{\mathbf{R}}_0^{-1}) \hat{\mathbf{r}}_\ell \quad (3)$$

where $\hat{\mathbf{r}}_\ell = \text{vec} \hat{\mathbf{R}}_\ell'$ is a $1 \times k^2$ row vector with rows of $\hat{\mathbf{R}}_\ell$ stacked one next to the other, and m is the lag order. The \otimes denotes the Kronecker product (http://en.wikipedia.org/wiki/Kronecker_product), $\hat{\mathbf{R}}_\ell = \mathbf{L}' \hat{\mathbf{\Gamma}}_\ell \mathbf{L}$, $\mathbf{L} \mathbf{L}' = \hat{\mathbf{\Gamma}}_0^{-1}$ where $\hat{\mathbf{\Gamma}}_\ell = n^{-1} \sum_{t=\ell+1}^n \hat{\mathbf{a}}_t \hat{\mathbf{a}}_{t-\ell}'$ is the lag ℓ residual autocovariance matrix.

The asymptotic distributions of \tilde{Q}_m is chi-squared with $k^2(m - p - q)$ degrees of freedom. In **portest** package, this statistic is implemented in the function `Hosking()`:

```
Hosking(obj, lags = seq(5, 30, 5), order = 0, SquaredQ = FALSE),
```

where the arguments of this function is described as before. Note that the function `portest()` with the arguments `test = "Hosking"`, `MonteCarlo = FALSE`, and `order = 0` will gives the same results of the function `Hosking()`. The Monte-Carlo version of this test statistic is implemented in the function `portest()` as an argument `test = "Hosking"` provided that `MonteCarlo = TRUE` is selected.

```
portest(obj, lags = seq(5, 30, 5), order = 0, test = "Hosking", MonteCarlo = TRUE,
        nslaves = 1, NREP = 1000, InfiniteVarianceQ = FALSE, SquaredQ = FALSE)
```

3.1. Example 4

In this example, we consider fitting a VAR(k), $k = 1, 2, 3$ model to the monthly log returns of the IBM stock and the S&P 500 index from January 1926 to December 1999 with 888 observations (Tsay 2005, p. 356). The p-values for the modified portmanteau test of Hosking (1980), \tilde{Q}_m , are computed using the Monte-Carlo test procedure with 10^3 replications. For additional comparisons, the p-values for \tilde{Q}_m are also evaluated using asymptotic approximations.

```
R> IBMSP500 <- monthibmspln
R> FitIBMSP5001 <- ar.ols(IBMSP500, aic = TRUE, intercept = F, order.max = 1)
R> Hosking(FitIBMSP5001)
```

Lags	Statistic	df	p-value
5	38.33044	16	0.0013574550
10	61.42150	36	0.0051949240
15	72.97170	56	0.0633819777
20	118.87159	76	0.0012179623
25	152.37966	96	0.0002208340
30	171.72563	116	0.0006001655

```
R> portest(FitIBMSP5001, test = "Hosking", nslaves = nslaves)
```

```
2 slaves are spawned successfully. 0 failed.
```

Lags	Statistic	df	p-value
5	38.33044	16	0.002997003
10	61.42150	36	0.003996004
15	72.97170	56	0.072927073
20	118.87159	76	0.003996004
25	152.37966	96	0.000999001
30	171.72563	116	0.002997003

```
R> FitIBMSP5002 <- ar.ols(IBMSP500, aic = TRUE, intercept = F, order.max = 2)
R> Hosking(FitIBMSP5002)
```

Lags	Statistic	df	p-value
5	28.12271	12	0.005307838
10	50.23144	32	0.021174563
15	61.53279	52	0.171676954
20	104.28887	72	0.007697842
25	138.24856	92	0.001303988
30	156.56512	112	0.003487092

```
R> portest(FitIBMSP5002, test = "Hosking", nslaves = nslaves)
```

```
2 slaves are spawned successfully. 0 failed.
```

Lags	Statistic	df	p-value
5	28.12271	12	0.003996004
10	50.23144	32	0.020979021
15	61.53279	52	0.145854146
20	104.28887	72	0.013986014
25	138.24856	92	0.001998002
30	156.56512	112	0.005994006

```
R> FitIBMSP5003 <- ar.ols(IBMSP500, aic = TRUE, intercept = F, order.max = 3)
R> Hosking(FitIBMSP5003)
```

Lags	Statistic	df	p-value
5	18.08797	8	0.020576519
10	40.78971	28	0.056135837
15	52.21967	48	0.313383239
20	93.82650	68	0.020716599
25	124.25318	88	0.006631765
30	142.81916	108	0.013972657

```
R> portest(FitIBMSP5003, test = "Hosking", nslaves = nslaves)
```

```

      2 slaves are spawned successfully. 0 failed.
Lags Statistic  df    p-value
  5   18.08797   8  0.029970030
 10   40.78971  28  0.053946054
 15   52.21967  48  0.307692308
 20   93.82650  68  0.023976024
 25  124.25318  88  0.004995005
 30  142.81916 108  0.018981019

```

All results reject the fitted VAR(1), VAR(2) and VAR(3) models.

3.2. Example 5

The trivariate quarterly time series, 1960–1982, of West German investment, income, and consumption was discussed by Lütkepohl (2005, §3.23). So $n = 92$ and $k = 3$ for this series. As in Lütkepohl (2005, §4.24) we model the logarithms of the first differences. Using the AIC and FPE, Lütkepohl (2005, Table 4.25) selected a VAR(2) for this data. All diagnostic tests reject simple randomness, VAR(0). The asymptotic distribution and the Monte-Carlo tests for VAR(1) suggests model inadequacy supports the choice of the VAR(2) model.

```

R> data("WestGerman")
R> DiffData <- matrix(numeric(3 * 91), ncol = 3)
R> for (i in 1:3) DiffData[, i] <- diff(log(WestGerman[, i]), lag = 1)
R> FitWestGerman <- ar.ols(DiffData, aic = FALSE, order.max = 2,
+   intercept = FALSE)
R> Hosking(FitWestGerman)

```

```

Lags Statistic  df    p-value
  5   30.36128  27  0.2981674
 10   71.94191  72  0.4797610
 15  122.49894 117  0.3455266
 20  171.96132 162  0.2811881
 25  209.45688 207  0.4391932
 30  254.48482 252  0.4443308

```

```

R> portest(FitWestGerman, test = "Hosking", nslaves = nslaves)

```

```

      2 slaves are spawned successfully. 0 failed.
Lags Statistic  df    p-value
  5   30.36128  27  0.3796204
 10   71.94191  72  0.5064935
 15  122.49894 117  0.3546454
 20  171.96132 162  0.2667333
 25  209.45688 207  0.4265734
 30  254.48482 252  0.4235764

```


4. Li and McLeod portmanteau test

Li and McLeod (1981) suggested the multivariate modified portmanteau test statistic

$$\tilde{Q}_m^{(L)} = n \sum_{\ell=1}^m \hat{\mathbf{r}}_\ell' (\hat{\mathbf{R}}_0^{-1} \otimes \hat{\mathbf{R}}_0^{-1}) \hat{\mathbf{r}}_\ell + \frac{k^2 m(m+1)}{2n} \quad (4)$$

which is distributed as chi-squared with $k^2(m-p-q)$ degrees of freedom. In **portes** package, the test statistic $\tilde{Q}_m^{(L)}$ is implemented in the function `LiMcLeod()`,

```
LiMcLeod(obj, lags = seq(5, 30, 5), order = 0, SquaredQ = FALSE),
```

where the arguments of this function is described as before.

Note that the function `portest()` with the arguments `test = "LiMcLeod"`, `MonteCarlo = FALSE`, and `order = 0` will gives the same results of the function `LiMcLeod()`. The Monte-Carlo version of this test statistic is implemented in the function `portest()` as an argument `test = "LiMcLeod"` provided that `MonteCarlo = TRUE` is selected.

```
portest(obj, lags = seq(5, 30, 5), order = 0, test = "LiMcLeod", MonteCarlo = TRUE,
        nslaves = 1, NREP = 1000, InfiniteVarianceQ = FALSE, SquaredQ = FALSE)
```

5. Generalized variance portmanteau test

Peña and Rodriguez (2002) proposed a univariate portmanteau test of goodness-of-fit test based on the m -th root of the determinant of the m -th Toeplitz residual autocorrelation matrix

$$\hat{\mathcal{R}}_m = \begin{pmatrix} \hat{r}_0 & \hat{r}_1 & \dots & \hat{r}_m \\ \hat{r}_{-1} & \hat{r}_0 & \dots & \hat{r}_{m-1} \\ \vdots & \dots & \ddots & \vdots \\ \hat{r}_{-m} & \hat{r}_{-m+1} & \dots & \hat{r}_0 \end{pmatrix} \quad (5)$$

where $\hat{r}_0 = 1$ and $\hat{r}_{-\ell} = \hat{r}_\ell$, for all ℓ . They approximated the distribution of their proposed test statistic by the gamma distribution and provided simulation experiments to demonstrate the improvement of their statistic in comparison with the one that is given in Eq. (2).

Peña and Rodriguez (2006) suggested to modify this test by taking the log of the $(m+1)$ -th root of the determinant in Eq. (5). They proposed two approximations by using the Gamma and Normal distributions to the asymptotic distribution of this test and indicated that the performance of both approximations for checking the goodness-of-fit in linear models is similar and more powerful for small sample size than the previous one. Lin and McLeod (2006) introduced the Monte-Carlo version of this test as they noted that it is quite often that the generalized variance portmanteau test does not agree with the suggested Gamma approximation. Mahdi and McLeod (2011) generalized both methods to the multivariate time series,

$$\mathfrak{D}_m = \frac{-3n}{2m+1} \log |\hat{\mathfrak{R}}_m|, \quad (6)$$

where

$$\hat{\mathfrak{R}}_m = \begin{pmatrix} \mathbb{I}_k & \hat{R}_1 & \dots & \hat{R}_m \\ \hat{R}_{-1} & \mathbb{I}_k & \dots & \hat{R}_{m-1} \\ \vdots & \dots & \ddots & \vdots \\ \hat{R}_{-m} & \hat{R}_{-m+1} & \dots & \mathbb{I}_k \end{pmatrix}. \quad (7)$$

The null distribution is approximately χ^2 with $k^2(1.5m(m+1)(2m+1)^{-1} - p - q)$ degrees of freedom and it is implemented in the R function `gvtest()`,

```
gvtest(obj, lags = seq(5, 30, 5), order = 0, SquaredQ = FALSE),
```

where the arguments of this function are described as before.

Note that the function `portest()` with the arguments `test = "gvtest"`, `MonteCarlo = FALSE`, and `order = 0` will give the same results of the function `gvtest()`. The Monte-Carlo version of this test statistic is implemented in the function `portest()` as an argument `test = "gvtest"` provided that `MonteCarlo = TRUE` is selected.

```
portest(obj, lags = seq(5, 30, 5), order = 0, test = "gvtest", MonteCarlo = TRUE,
        nslaves = 1, NREP = 1000, InfiniteVarianceQ = FALSE, SquaredQ = FALSE)
```

5.1. Example 6

Consider again the log numbers of Canadian lynx trappings univariate series from 1821 to 1934, where the AR(2) model is selected based on the BIC criterion using the function `SelectModel` in the R package **FitAR** as a first step in the analysis. Now, we apply the statistic \mathfrak{D}_m on the fitted model based on the asymptotic distribution and the Monte-Carlo significance test,

```
R> gvtest(Fitlynx)
```

Lags	Statistic	df	p-value
5	5.984989	2.090909	0.054687987
10	10.036630	5.857143	0.115222212
15	21.447021	9.612903	0.014964682
20	31.810564	13.365854	0.003100578
25	38.761595	17.117647	0.002040281
30	43.936953	20.868852	0.002252062

```
R> portest(Fitlynx, test = "gvtest", nslaves = nslaves)
```

2 slaves are spawned successfully. 0 failed.

Lags	Statistic	df	p-value
5	5.984989	2.090909	0.063936064
10	10.036630	5.857143	0.080919081
15	21.447021	9.612903	0.006993007
20	31.810564	13.365854	0.002997003
25	38.761595	17.117647	0.000999001
30	43.936953	20.868852	0.000999001

After that, we fit the subset autoregressive AR_(1,2,4,10,11) using the BIC and then we apply \mathfrak{D}_m as before,

```
R> gvtest(FitsubsetAR)
```

Lags	Statistic	df	p-value
5	2.374225	0.0000000	NA
10	3.598248	0.0000000	NA
15	5.661285	0.6129032	0.008190694
20	8.590962	4.3658537	0.090004731
25	11.462473	8.1176471	0.184353957
30	13.900470	11.8688525	0.297764350

```
R> portest(FitsubsetAR, test = "gvtest", nslaves = nslaves)
```

2 slaves are spawned successfully. 0 failed.

Lags	Statistic	df	p-value
5	2.374225	0.0000000	0.3476523
10	3.598248	0.0000000	0.6703297
15	5.661285	0.6129032	0.7152847
20	8.590962	4.3658537	0.6953047
25	11.462473	8.1176471	0.6663337
30	13.900470	11.8688525	0.6793207

However the approximation asymptotic distribution of the statistic \mathfrak{D}_m suggests that the subset AR model is an adequate model for lags $m \geq 20$, the Monte-Carlo portmanteau test is clearly suggest that the subset AR model is an adequate model.

5.2. Example 7

consider again fitting a VAR(k), $k = 1, 2, 3$ model to the monthly log returns of the IBM stock and the S&P 500 index from January 1926 to December 1999 with 888 observations ([Tsay 2005](#), p. 356).

```
R> gvtest(FitIBMSP5001)
```

Lags	Statistic	df	p-value
5	26.73298	12.36364	0.010069145
10	50.16580	27.42857	0.005076554
15	66.95921	42.45161	0.009606334
20	87.59443	57.46341	0.006384252
25	108.82328	72.47059	0.003699716
30	128.30068	87.47541	0.002940363

```
R> portest(FitIBMSP5001, test = "gvtest", nslaves = nslaves)
```

2 slaves are spawned successfully. 0 failed.

Lags	Statistic	df	p-value
5	26.73298	12.36364	0.002997003
10	50.16580	27.42857	0.003996004
15	66.95921	42.45161	0.006993007
20	87.59443	57.46341	0.004995005
25	108.82328	72.47059	0.005994006
30	128.30068	87.47541	0.003996004

R> *gvtest*(FitIBMSP5002)

Lags	Statistic	df	p-value
5	16.24518	8.363636	0.04647938
10	38.00564	23.428571	0.02910435
15	54.26122	38.451613	0.04688480
20	74.35787	53.463415	0.03091912
25	95.55057	68.470588	0.01701104
30	114.96754	83.475410	0.01272103

R> *portest*(FitIBMSP5002, test = "gvtest", nslaves = nslaves)

2 slaves are spawned successfully. 0 failed.

Lags	Statistic	df	p-value
5	16.24518	8.363636	0.006993007
10	38.00564	23.428571	0.007992008
15	54.26122	38.451613	0.016983017
20	74.35787	53.463415	0.017982018
25	95.55057	68.470588	0.008991009
30	114.96754	83.475410	0.008991009

R> *gvtest*(FitIBMSP5003)

Lags	Statistic	df	p-value
5	6.914649	4.363636	0.16977954
10	24.655501	19.428571	0.18989321
15	39.324113	34.451613	0.26078729
20	58.297021	49.463415	0.18238250
25	79.102500	64.470588	0.10384117
30	98.361967	79.475410	0.07416598

R> *portest*(FitIBMSP5003, test = "gvtest", nslaves = nslaves)

2 slaves are spawned successfully. 0 failed.

Lags	Statistic	df	p-value
5	6.914649	4.363636	0.05394605
10	24.655501	19.428571	0.06493506

```

15 39.324113 34.451613 0.11488511
20 58.297021 49.463415 0.08891109
25 79.102500 64.470588 0.06093906
30 98.361967 79.475410 0.04995005

```

While the fitted VAR (1) and VAR (2) models are rejected, the \mathfrak{D}_m diagnostic suggests that the fitted VAR (3) maybe consider to be an adequate model.

5.3. Example 8

In this example, we consider the quarterly time series, 1960–1982, of West German investment, income, and consumption studied before.

We apply the statistic \mathfrak{D}_m on the fitted VAR (2) model based on the asymptotic distribution and the Monte-Carlo significance test,

```
R> gvtest(FitWestGerman)
```

Lags	Statistic	df	p-value
5	20.90960	18.81818	0.3310522834
10	52.17337	52.71429	0.4951413528
15	91.80348	86.51613	0.3283404972
20	135.40962	120.29268	0.1637652676
25	195.17389	154.05882	0.0139543395
30	257.76048	187.81967	0.0005343724

```
R> portest(FitWestGerman, test = "gvtest", nslaves = nslaves)
```

```

      2 slaves are spawned successfully. 0 failed.
Lags Statistic      df    p-value
  5  20.90960   18.81818 0.3116883
 10  52.17337   52.71429 0.5424575
 15  91.80348   86.51613 0.5624376
 20 135.40962  120.29268 0.5954046
 25 195.17389  154.05882 0.4005994
 30 257.76048  187.81967 0.3486513

```

Using the asymptotic distribution, results suggest that the VAR (2) model is adequate at lags $m < 25$ and inadequate at lags $m \geq 25$, whereas Monte-Carlo test supports the choice of the VAR (2) model.

References

Box G, Pierce D (1970). “Distribution of Residual Autocorrelation in Autoregressive-Integrated Moving Average Time Series Models.” *Journal of American Statistical Association*, **65**, 1509–1526.

- Hosking JRM (1980). “The Multivariate Portmanteau Statistic.” *Journal of American Statistical Association*, **75**(371), 602–607.
- Li WK, McLeod AI (1981). “Distribution of the Residual Autocorrelation in Multivariate ARMA Time Series Models.” *Journal of the Royal Statistical Society, Series B*, **43**(2), 231–239.
- Lin J, McLeod AI (2006). “Improved Peña-Rodriguez Portmanteau Test.” *Computational Statistics and Data Analysis*, **51**(3), 1731–1738.
- Ljung G, Box G (1978). “On a Measure of Lack of Fit in Time Series Models.” *Biometrika*, **65**, 297–303.
- Lütkepohl H (2005). *New Introduction to Multiple Time Series Analysis*. Springer-Verlag, New York.
- Mahdi E, McLeod AI (2011). “Improved Multivariate Portmanteau Diagnostic Test.” Submitted.
- McLeod A, Zhang Y (2008). “Improved Subset Autoregression: With R Package.” *Journal of Statistical Software*, **28**(2), 1–28. URL <http://www.jstatsoft.org/v28/i02>.
- Peña D, Rodriguez J (2002). “A Powerful Portmanteau Test of Lack of Test for Time Series.” *Journal of American Statistical Association*, **97**(458), 601–610.
- Peña D, Rodriguez J (2006). “The Log of the Determinant of the Autocorrelation Matrix for Testing Goodness of Fit in Time Series.” *Journal of Statistical Planning and Inference*, **8**(136), 2706–2718.
- Tsay RS (2005). *Analysis of Financial Time Series*. Wiley, New York, 2nd edition.

Affiliation:

A. Ian McLeod
Department of Statistical and Actuarial Sciences
University of Western Ontario
E-mail: aim@stats.uwo.ca
URL: <http://www.stats.uwo.ca/mcleod>
Esam Mahdi
Department of Statistical and Actuarial Sciences
University of Western Ontario
E-mail: emahdi@uwo.ca